

COVID-19 and ABO blood group: another viewpoint

Li *et al.*¹ have recently published 'Association between ABO blood groups and risk of SARS-CoV-2 pneumonia', an observation already reported a few weeks ago as a MedRxiv preprint by Zhao *et al.*² and which had a certain impact in the press.

In both studies, the ABO blood groups distribution of patients with coronavirus disease 2019 (COVID-19) were compared to that of controls from the local populations that showed that blood group A was associated with an increased risk of infection, whereas group O was associated with a decreased risk. Considering this information rather as a working hypothesis, some scientists have called for caution.³

However, as already strongly suggested by others,⁴ this variable susceptibility to severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection could be linked to circulating anti-A antibodies, which could interfere or even inhibit the virus–cell adhesion process.

We had the idea to analyse these important available data series from the anti-A or -B antibodies viewpoint instead of ABO blood group antigens as the authors did.

In fact, considering the largest series of patients with COVID-19 ($N = 1888$) analysed by Zhao *et al.*,² we compared the proportion of those possessing anti-A in their serum (i.e. those of B and O blood groups) and those who did not (i.e. those of A and AB blood groups) to the control cohort ($N = 3694$; Table I).

The results (Table I) indicate that subjects with anti-A in serum (i.e. B and O blood groups) are significantly less represented in the COVID-19 group than those lacking anti-A

whatever the group: A and AB ($P < 0.001$), A ($P < 0.001$) or AB ($P = 0.0323$), whereas there was no significant difference versus circulating anti-B (data not shown).

We then wondered if there was a difference between anti-A from O and anti-A from B, and then we compared the supposed protective effect of anti-A from O and from B (Table II).

Whereas both blood group O and B patients possess circulating seric anti-A, it appears and it is statistically highly significant ($P < 0.001$) that O group patients are underrepresented (49.4 % vs. 57.6%), whereas B group patients are, on the contrary, overrepresented (50.6% vs. 42.4%), meaning that anti-A from O is more protective than anti-A from B.

This latter observation is probably related to the fact that the immunoglobulin predominant isotype of anti-B/anti-A is IgM in serum from group A and B individuals, but IgG in O group serum, an already known notion,⁵ which has been well documented by flow cytometry.⁶

In conclusion, this way of analysing the data strongly suggests that the presence of anti-A antibodies in serum and more specifically IgG anti-A, should be considered as a factor more significant than the blood group itself, as far as the relationship between COVID-19 susceptibility and ABO blood groups is concerned.

Far from intending to corroborate the authors' conclusions as such, we wanted to show that the resources of immuno-haematology allow several approaches that could perhaps be useful for the disease follow-up.

Table I. Comparison of subjects with/without anti-A antibodies in their serum.

	RBC blood group	Control, n (%)	COVID-19, n (%)	χ^2	P	OR (95% CI)
With anti-A	B and O	2170 (58.7)	927 (52.2)			
	A and AB	1524 (41.3)	848 (47.8)	20.74	<0.001	1.30 (1.16–1.46)
Without anti-A	A	1188 (32.2)	670 (37.7)	19.97	<0.001	1.32 (1.17–1.49)
	AB	336 (9.1)	178 (10.0)	4.58	0.0323	1.24 (1.02–1.51)

Table II. Comparison of anti-A from O and from B subjects.

	RBC blood group	Control, n (%)	COVID-19, n (%)	χ^2	P	OR (95% CI)
Anti-A from O	O	1250 (57.6)	458 (49.4)			
Anti-A from B	B	920 (42.4)	469 (50.6)	17.64	<0.001	1.39 (1.19–1.62)

Christiane Gérard¹ 
Gianni Maggipinto¹ 
Jean-Marc Minon² 

¹Blood Transfusion service, Centre Hospitalier Universitaire, University of Liège, Liège, Belgium and ²Department of Thrombosis-haemostasis and Transfusion, Centre Hospitalier Régional de la citadelle, Liège, Belgium.
E-mail: gerardchristiane4@gmail.com

Keywords: antibodies, blood group serology, COVID-19

References

1. Li J, Wang X, Chen J, Cai Y, Deng A, Yang M. Association between ABO blood groups and risk of SARS-CoV-2 pneumonia. *Br J Haematol*. 2020 [Epub ahead of print]. DOI: <https://doi.org/10.1111/bjh.16797>.
2. Zhao J, Yang Y, Huang H, Li D, Gu D, Lu X, et al. Relationship between the ABO Blood Group and the COVID-19 Susceptibility. *MedRxiv* 2020 [Epub ahead of print]. DOI: <https://doi.org/10.1101/2020.03.11.20031096>.
3. Pirenne F, Chiaroni J. Que penser des données sur la susceptibilité moindre des individus O à l'infection au COVID 19? 2020. Available at: https://www.sfts.asso.fr/Media/20200411_abo_et_covid_19_-_f.pirenne_j.chiaroni.pdf. Accessed May 2020
4. Guillon P, Clément M, Sébille V, Rivain J-G, Chou C-F, Ruvoën-Clouet N, et al. Inhibition of the interaction between the SARS-CoV Spike protein and its cellular receptor by anti-histo-blood group antibodies. *Glycobiology*. 2008;**18**:1085–93.
5. Fung, MK, Grossman, BJ, Hiller, CD & Westhoff, CM, eds. *American Association of Blood Banks (AABB) Technical Manual*, 18th edn. Washington, DC: American Association of Blood Banks, 2014: 297.
6. Stussi G, Huggel K, Lutz HU, Schanz U, Rieben R, Seebach JD. Isotype-specific detection of ABO blood group antibodies using a novel flow cytometric method. *Br J Haematol*. 2005;**130**:954–63.